

January 1999
 hep-th/9902001

Algebraic Structure in $0 < c \leq 1$ Open-Closed String Field Theories

Daiji Ennyu¹

Hiroshima National College of Maritime Technology

Toyota-Gun, Hiroshima 725-0200, Japan

Hiroshi Kawabe²

Yonago National College of Technology

Yonago 683-8502, Japan

and

Naohito Nakazawa³

Department of Physics, Faculty of Science and Engineering, Shimane University

Matsue 690-8504, Japan

Abstract

We apply stochastic quantization method to Kostov's matrix-vector models for the second quantization of orientable strings with Chan-Paton like factors, including both open and closed strings. The Fokker-Planck hamiltonian deduces an orientable open-closed string field theory at the double scaling limit. There appears an algebraic structure in the continuum F-P hamiltonian including a Virasoro algebra and a $SU(r)$ current algebra.

¹e-mail address: ennyu@hiroshima-cmt.ac.jp

²e-mail address: kawabe@yonago-k.ac.jp

³e-mail address: nakazawa@ifse1.riko.shimane-u.ac.jp

The explicit construction of non-critical string field theories via the double scaling limit of the matrix models [1] provides not only the basis for the non-perturbative analysis of string theories but also the clear understanding of the origin of the constraints realized in the algebraic structure of the string field theoretic hamiltonian [1-10]. The algebraic structure in non-critical string field theories has been investigated in this context for orientable closed strings [2, 3, 4], non-orientable closed strings [6], orientable open-closed strings [7, 8] and non-orientable open-closed strings[10]. For orientable 2D surfaces with boundaries, an interesting algebraic structure, $sl(r, \mathbf{C}) \times sl(r, \mathbf{C})$ chiral current algebra including $SU(r)$ current algebra, has been found in a field theoretic hamiltonian of orientable open-closed strings[9]. The observation indicates the relation between the algebraic structure and a Chan-Paton like realization of gauge groups in open string theories. While the $SO(r)$ current algebra has been found in the string field theoretic hamiltonian for non-orientable open-closed strings[10].

In this note, we construct non-critical orientable open-closed string field theories for $0 < c \leq 1$ by applying stochastic quantization method to Kostov's matrix-vector model[7, 11]. To introduce Chan-Paton like factors, we slightly modify Kostov's model. The continuum limit of the string field theory is taken at the double scaling limit of this matrix-vector model. We introduce the scale parameters on the boundaries which characterize the scaling behaviour of the time evolution along the boundaries. We show that the string field theoretic hamiltonian, which is the continuum limit of the loop space Fokker-Planck hamiltonian in stochastic quantization[3, 6], appears in the form of a linear combination of three deformation generators of orientable strings which satisfy algebraic relations including a Virasoro algebra and a $SU(r)$ current algebra.

For orientable 2D surfaces with boundaries, the sum of triangulated surfaces is reconstructed by the hermitian matrix-vector models[11, 12]. To introduce the orientable open-closed string interactions for $0 < c \leq 1$, we consider Kostov's matrix-vector model with Chan-Paton like factors. The action is given by[7]

$$S_1 = \text{Tr} \left(\frac{1}{2} \sum_{x,x'} A_{xx'} A_{x'x} - \frac{1}{3} \frac{g}{\sqrt{N}} \sum_{x,x',x''} A_{xx'} A_{x'x''} A_{x''x} \right),$$

$$S_2 = \sum_{x,x'} \sum_{a=1}^r V_x^{a*} (\delta_{xx'} - \frac{g_B^a}{\sqrt{N}} A_{xx'}) V_{x'}^a, \quad (1)$$

where the interactions of matrices, $A_{xx'}$, take place only for $|x-x'| \leq 1$ in the target space $\mathbf{Z}$, i.e. $x, x' \in \mathbf{Z}$. Namely, the relevant matrices are $A_{xx \ ij}$, which are $N \times N$ hermitian matrices, and $A_{xx+1 \ ij} = A_{x+1x \ ij}^\dagger$. $V_{x \ i}^a$ and $V_{x \ i}^{a*}$ are N -dimensional complex vectors with a Chan-Paton like factor a which runs from 1 to r . The partition function, Z , and the free energy, F , of this model are given by

$$\begin{aligned} Z &\equiv \int dA dV dV^* e^{-S_1 - S_2}, \\ F &\equiv \ln Z \\ &= \prod_{a=1}^r (g_B^a)^{L_B^a} g^S N^\chi, \end{aligned} \quad (2)$$

where S and L_B^a are the total area and the total length of the boundary with the “colour” index a of the 2D surface, respectively. χ is the Euler number of the triangulated 2D surface with boundaries, $\chi \equiv 2 - 2\#(\text{handles}) - \#(\text{boundaries})$. By integrating out all the non-hermitian matrices, $A_{xx+1 \ ij}$ and $A_{x+1x \ ji}^\dagger$, we obtain an effective action as a function of the hermitian matrix $M_{x \ ij} \equiv A_{xx \ ij} - \frac{\sqrt{N}}{2g} \delta_{ij}$, and the vectors, $V_{x \ i}^a$ and $V_{x \ i}^{a*}$. We denote the effective action as $S_{\text{eff}}(M, V, V^*)$ [7].

We define the time evolution, $M_{x \ ij}(\tau + \Delta\tau) \equiv M_{x \ ij}(\tau) + \Delta M_{x \ ij}(\tau)$ and $V_{x \ i}^a(\tau + \Delta\tau) \equiv V_{x \ i}^a(\tau) + \Delta V_{x \ i}^a(\tau)$, in terms of Ito’s calculus[13] with respect to the stochastic time τ which is discretized with the unit time step $\Delta\tau$. The time evolution is given by the following Langevin equations,

$$\begin{aligned} \Delta M_{x \ ij}(\tau) &= -\frac{\partial}{\partial M_{x \ ji}} S_{\text{eff}}(M, V, V^*)(\tau) \Delta\tau + \Delta \xi_{x \ ij}(\tau), \\ \Delta V_{x \ i}^a(\tau) &= -\lambda_x^a \frac{\partial}{\partial V_{x \ i}^{a*}} S_{\text{eff}}(M, V, V^*)(\tau) \Delta\tau + \Delta \eta_{x \ i}^a(\tau), \\ \Delta V_{x \ i}^{a*}(\tau) &= -\lambda_x^a \frac{\partial}{\partial V_{x \ i}^a} S_{\text{eff}}(M, V, V^*)(\tau) \Delta\tau + \Delta \eta_{x \ i}^{a*}(\tau). \end{aligned} \quad (3)$$

We have introduced the scale parameter λ_x^a in the Langevin equation of vector variables, $V_{x \ i}^a$ and $V_{x \ i}^{a*}$. At the double scaling limit, the parameter λ_x^a defines the time scale for the stochastic time evolution of the open string end-point along the boundaries with the “colour” index a .

The correlations of the white noise $\Delta\xi_{x\ ij}$ and $\Delta\eta_{x\ i}^a$ are defined by

$$\begin{aligned} < \Delta\xi_{x\ ij}(\tau)\Delta\xi_{x'\ kl}(\tau) >_{\xi} &= 2\Delta\tau\delta_{il}\delta_{jk}\delta_{xx'} , \\ < \Delta\eta_{x\ i}^{a*}(\tau)\Delta\eta_{x'\ j}^b(\tau) >_{\eta} &= 2\lambda_x^a\Delta\tau\delta^{ab}\delta_{ij}\delta_{xx'} . \end{aligned} \quad (4)$$

We will see later that the λ_x^a -independence of the *equilibrium* is realized as the algebraic relation between the constraints in the string field theoretic hamiltonian which is equivalent to the integrability condition of the multi-time evolution of the Langevin equations.

Now we define the closed strings, $\phi_x(L) = N^{-1}\text{tr}(e^{LN^{-1/2}M_x})$, and the open strings, $\psi_x^{ab}(L) = N^{-1}(V^{a*}e^{LN^{-1/2}M_x}V^b)$. Following to Ito's calculus, we calculate the time development of these string variables[6],

$$\begin{aligned} \Delta\phi_x(L) &= \Delta\tau L\left\{ \int_0^L dL' \phi_x(L')\phi_x(L-L') + \sum_{x'} \int_0^\infty dL' C_{xx'}^{(c)} \phi_x(L+L')\phi_{x'}(L') \right. \\ &+ \frac{1}{Ng} \sum_{a,b,x'} g_B^a g_B^b \int_0^\infty dL' L' C_{xx'}^{(o)} \psi_x^{ab}(L+L')\psi_{x'}^{ba}(L') \\ &+ \left. \frac{1}{N} \sum_{a=1}^r g_B^a \psi_x^{aa}(L) - \frac{1}{4g} \phi_x(L) + g \frac{\partial^2}{\partial L^2} \phi_x(L) \right\} + \Delta\zeta_x(L) , \\ \Delta\psi_x^{ab}(L) &= \lambda_x^a \Delta\tau \left\{ \left(-1 + \frac{g_B^a}{2g} + g_B^a \frac{\partial}{\partial L}\right) \psi_x^{ab}(L) + \frac{1}{g} \sum_{c,x'} g_B^a g_B^c \int_0^\infty dL' C_{xx'}^{(o)} \psi_x^{cb}(L+L')\psi_{x'}^{ac}(L') \right\} \\ &+ \lambda_x^b \Delta\tau \left\{ \left(-1 + \frac{g_B^b}{2g} + g_B^b \frac{\partial}{\partial L}\right) \psi_x^{ab}(L) + \frac{1}{g} \sum_{c,x'} g_B^b g_B^c \int_0^\infty dL' C_{xx'}^{(o)} \psi_x^{ac}(L+L')\psi_{x'}^{cb}(L') \right\} \\ &+ 2\Delta\tau \lambda_x^a \delta^{ab} \phi_x(L) \\ &+ \Delta\tau \left\{ -\frac{L}{4g} \psi_x^{ab}(L) + Lg \frac{\partial^2}{\partial L^2} \psi_x^{ab}(L) + L \sum_{x'} \int_0^\infty dL' C_{xx'}^{(c)} \psi_x^{ab}(L+L')\phi_{x'}(L') \right. \\ &+ \sum_c g_B^c \int_0^L dL' \psi_x^{ac}(L')\psi_x^{cb}(L-L') \\ &+ \sum_{c,d,x'} \frac{g_B^c g_B^d}{g} \int_0^L dL' \int_0^\infty dL'' \int_0^\infty dL''' C_{xx'}^{(o)} \psi_x^{ad}(L'+L'')\psi_x^{cb}(L-L'+L''')\psi_{x'}^{dc}(L''+L''') \\ &+ \left. 2 \int_0^L dL' L' \psi_x^{ab}(L')\phi_x(L-L') \right\} + \Delta\zeta_x^{ab}(L) . \end{aligned} \quad (5)$$

Here we have introduced the adjacency matrices, $C_{xx'}^{(c)} = C_{xx'}^{(o)} = \delta_{xx'+1} + \delta_{xx'-1}$. In (5), the new noise variables are defined by

$$\Delta\zeta_x(L) \equiv LN^{-3/2}\text{tr}(e^{LN^{-1/2}M_x}\Delta\xi_x) ,$$

$$\begin{aligned}\Delta\zeta_x^{ab}(L) &\equiv N^{-3/2} \int_0^L dL' V_x^{a*} e^{L'N^{-1/2}M_x} \Delta\xi_x e^{(L-L')N^{-1/2}M_x} V_x^b \\ &+ N^{-1} \{ V_x^{a*} e^{LN^{-1/2}M_x} \Delta\eta_x^b + \Delta\eta_x^{a*} e^{LN^{-1/2}M_x} V_x^b \} .\end{aligned}\quad (6)$$

We notice that $\langle \Delta\zeta_x(L) \rangle_{\xi\eta} = \langle \Delta\zeta_x^{ab}(L) \rangle_{\xi\eta} = 0$ hold in the sense of Ito's calculus. Especially, from eqs. (5), the λ -independence of the equilibrium limit deduces a S-D equation which ensures the $SU(r)$ current algebra.

We can describe a class of matrix-vector models for the central charge $0 < c \leq 1$ with the following adjacency matrices[7].

$$\begin{aligned}C_{xx'}^{(c)} &= \cos(\pi p_0)(\delta_{xx'+1} + \delta_{xx'-1}) , \\ C_{xx'}^{(o)} &= \cos(\pi p_0/2)(\delta_{xx'+1} + \delta_{xx'-1}) .\end{aligned}\quad (7)$$

The case, $p_0 = 1/m$, corresponds to the central charge, $c = 1 - \frac{6}{m(m+1)}$.

The stochastic process is interpreted as the time evolution in a string field theory. The corresponding non-critical orientable open-closed string field theory is defined by the F-P hamiltonian operator. In terms of the expectation value of an observable $O(\phi, \psi)$, a function of $\phi_x(L)$'s and $\psi_x^{ab}(L)$'s, the F-P hamiltonian operator $\hat{H}_{FP}$ is defined by[6]

$$\langle \phi(0), \psi(0) | e^{-\tau \hat{H}_{FP}} O(\hat{\phi}, \hat{\psi}) | 0 \rangle \equiv \langle O(\phi_{\xi\eta}(\tau), \psi_{\xi\eta}(\tau)) \rangle_{\xi\eta} . \quad (8)$$

In R.H.S., $\phi_{\xi\eta}(\tau)$ and $\psi_{\xi\eta}(\tau)$ denote the solutions of the Langevin equations (5) with the appropriate initial configuration $\phi_x(L; \tau = 0)$ and $\psi_x^{ab}(L; \tau = 0)$. In L.H.S., $\hat{H}_{FP}$ is given by the differential operator in the well-known Fokker-Planck equation for the expectation value of the observable $O(\phi, \psi)$ by replacing the closed (open) string variable $\phi_x(L)$ ($\psi_x^{ab}(L)$) to the creation operator $\hat{\phi}_x(L)$ ($\hat{\psi}_x^{ab}(L)$) and the differential $\frac{\partial}{\partial \phi_x(L)}$ ($\frac{\partial}{\partial \psi_x^{ab}(L)}$) to the annihilation operator $\hat{\pi}_x(L)$ ($\hat{\pi}_x^{ab}(L)$), respectively.

The continuum limit of $\hat{H}_{FP}$ is taken by introducing a length scale “ ϵ ” which defines the physical length of the strings as $l \equiv L\epsilon$. At the double scaling limit $\epsilon \rightarrow 0$, we keep the string coupling constant, $G \equiv N^{-2}\epsilon^{-2D}$, to be finite. The continuum stochastic time is given by $d\tau \equiv \epsilon^{-2+D}\Delta\tau$. While we define the scaling of the scale parameter λ_x^a by $\lambda^a \equiv \epsilon^{-1/2-D/2}\lambda_x^a$. We also redefine field variables as follows.

$$\Phi_x(l) \equiv \epsilon^{-D}\hat{\phi}_x(L) ,$$

$$\begin{aligned}
\Pi_x(l) &\equiv \epsilon^{-1+D} \hat{\pi}_x(L) , \\
\Psi_x^{ab}(l) &\equiv \epsilon^{-1/2-D/2} \hat{\psi}_x^{ab}(L) , \\
\Pi_x^{ab}(l) &\equiv \epsilon^{-1/2+D/2} \hat{\pi}_x^{ab}(L) .
\end{aligned} \tag{9}$$

The commutation relations are $[\Pi_x(l), \Phi_{x'}(l')] = \delta_{xx'} \delta(l-l')$ and $[\Pi_x^{ab}(l), \Phi_{x'}^{cd}(l')] = \delta^{ac} \delta^{bd} \delta_{xx'} \delta(l-l')$. Then we obtain the continuum F-P hamiltonian, $\mathcal{H}_{FP}$, from $\hat{H}_{FP}$,

$$\begin{aligned}
\mathcal{H}_{FP} = & \sum_x \int_0^\infty dl l \{ \sum_{x'} C_{xx'} \int_0^\infty dl' \Phi_x(l+l') \Phi_{x'}(l') + \sqrt{G} \sum_{a,b,x'} C_{xx'} \int_0^\infty dl' l' \Psi_x^{ab}(l+l') \Psi_{x'}^{ba}(l') \\
& + \int_0^l dl' \Phi_x(l') \Phi_x(l-l') + G \int_0^\infty dl' l' \Phi_x(l+l') \Pi_x(l') \} \Pi_x(l) \\
& + \sum_{a,b,x} \int_0^\infty dl \{ l \sum_{x'} C_{xx'} \int_0^\infty dl' \Psi_x^{ab}(l+l') \Phi_{x'}(l') \\
& + \sum_{c,d,x'} C_{xx'} \int_0^l dl_1 \int_0^\infty dl_2 \int_0^\infty dl_3 \Psi_x^{ad}(l_1+l_2) \Psi_x^{cb}(l-l_1+l_3) \Psi_{x'}^{dc}(l_2+l_3) \\
& + 2 \int_0^l dl' l' \Psi_x^{ab}(l') \Phi_x(l-l') \\
& + \lambda^a \sum_{c,x'} C_{xx'} \int_0^\infty dl' \Psi_x^{cb}(l+l') \Psi_{x'}^{ac}(l') + \lambda^b \sum_{c,x'} C_{xx'} \int_0^\infty dl' \Psi_x^{ac}(l+l') \Psi_{x'}^{cb}(l') \\
& + (\lambda^a + \lambda^b) \delta^{ab} \Phi_x(l) \} \Pi_x^{ab}(l) \\
& + \sum_{a,b,c,d,x} \sqrt{G} \int_0^\infty dl \int_0^\infty dl' \int_0^l dl_1 \int_0^{l'} dl'_1 \Psi_x^{ad}(l_1+l'_1) \Psi_x^{cb}(l+l'-l_1-l'_1) \Pi_x^{ab}(l) \Pi_x^{cd}(l') \\
& + \sum_{a,b,c,x} \sqrt{G} \int_0^\infty dl \int_0^\infty dl' \{ \lambda^a \Psi_x^{cb}(l+l') \Pi_x^{ab}(l) \Pi_x^{ca}(l') + \lambda^b \Psi_x^{ac}(l+l') \Pi_x^{ab}(l) \Pi_x^{bc}(l') \} \\
& + 2G \sum_{a,b,x} \int_0^\infty dl \int_0^\infty dl' l l' \Psi_x^{ab}(l+l') \Pi_x(l) \Pi_x^{ab}(l') .
\end{aligned} \tag{10}$$

The continuum F-P hamiltonian (10) takes the form of a linear combination of three continuum generators of string deformation,

$$\begin{aligned}
\mathcal{H}_{FP} = & \sum_x \int_0^\infty dl l \mathcal{L}_x(l) \Pi_x(l) + \sum_{a,b,x} \int_0^\infty dl \{ \lambda^a \mathcal{J}_x^{ab}(l) + \lambda^b \mathcal{J}_x^{ba*}(l) \} \Pi_x^{ab}(l) \\
& + \sum_{a,b,x} \int_0^\infty dl \{ l \mathcal{K}_x^{ab}(l) - \sum_c \int_0^l dl' l' (\mathcal{J}_x^{cb}(l') \Psi_x^{ac}(l-l') \\
& + \mathcal{J}_x^{ca*}(l') \Psi_x^{bc*}(l-l')) \} \Pi_x^{ab}(l) .
\end{aligned} \tag{11}$$

The explicit forms of these generators are given by

$$\mathcal{L}_x(l) = \sum_{x'} C_{xx'} \int_0^\infty dl' \Phi_x(l+l') \Phi_{x'}(l')$$

$$\begin{aligned}
& + \sqrt{G} \sum_{a,b,x'} C_{xx'} \int_0^\infty dl' l' \Psi_x^{ab}(l+l') \Psi_{x'}^{ba}(l') + \int_0^l dl' \Phi_x(l') \Phi_x(l-l') \\
& + G \int_0^\infty dl' l' \Phi_x(l+l') \Pi_x(l') + G \sum_{a,b} \int_0^\infty dl' l' \Psi_x^{ab}(l+l') \Pi_x^{ab}(l') , \\
\mathcal{K}_x^{ab}(l) & = \sum_{x'} C_{xx'} \int_0^\infty dl' \Psi_x^{ab}(l+l') \Phi_{x'}(l') \\
& + G \int_0^\infty dl' l' \Psi_x^{ab}(l+l') \Pi_x(l') + \sum_{c,d,x'} C_{xx'} \int_0^\infty dl_1 \int_0^{l+l_1} dl_2 \Psi_x^{ac}(l_2) \Psi_x^{db}(l+l_1-l_2) \Psi_{x'}^{cd}(l_1) \\
& + \sqrt{G} \sum_{c,d} \int_0^\infty dl_1 \int_0^{l+l_1} dl_2 \Psi_x^{ac}(l_2) \Psi_x^{db}(l+l_1-l_2) \Pi_x^{dc}(l_1) \\
& + 2 \int_0^l dl' \Psi_x^{ab}(l') \Phi_x(l-l') , \\
\mathcal{J}_x^{ab}(l) & = \delta^{ab} \Phi_x(l) + \sum_{c,x'} C_{xx'} \int_0^\infty dl' \Psi_x^{cb}(l+l') \Psi_{x'}^{ac}(l') \\
& + \sqrt{G} \sum_c \int_0^\infty dl' \Psi_x^{cb}(l+l') \Pi_x^{ca}(l') . \tag{12}
\end{aligned}$$

These generators satisfy the following algebra including the Virasoro algebra and the $SU(r)$ current algebra,

$$\begin{aligned}
[\mathcal{L}_x(l), \mathcal{L}_{x'}(l')] & = -(l-l') G \delta_{xx'} \mathcal{L}_x(l+l') , \\
[\mathcal{L}_x(l), \mathcal{K}_{x'}^{ab}(l')] & = -(l-l') G \delta_{xx'} \mathcal{K}_x^{ab}(l+l') \\
& + G \delta_{xx'} \sum_c \int_0^l du (l-u) \{ \mathcal{J}_x^{cb}(l+l'-u) \Psi_x^{ac}(u) + \mathcal{J}_x^{ca*}(l+l'-u) \Psi_x^{bc*}(u) \} , \\
[\mathcal{L}_x(l), \mathcal{J}_{x'}^{ab}(l')] & = l' G \delta_{xx'} \mathcal{J}_x^{ab}(l+l') , \\
[\mathcal{J}_x^{ab}(l), \mathcal{K}_{x'}^{cd}(l')] & = \sqrt{G} \delta^{ad} \delta_{xx'} \mathcal{K}_x^{cb}(l+l') \\
& - \sqrt{G} \delta^{ad} \delta_{xx'} \sum_e \int_0^l du \mathcal{J}_x^{ec*}(l+l'-u) \Psi_x^{be*}(u) \\
& - \sqrt{G} \delta_{xx'} \int_0^l du \mathcal{J}_x^{ad}(l+l'-u) \Psi_x^{cb}(u) , \\
[\mathcal{J}_x^{ab}(l), \mathcal{J}_{x'}^{cd}(l')] & = \sqrt{G} \delta^{ad} \delta_{xx'} \mathcal{J}_x^{cb}(l+l') - \sqrt{G} \delta^{cb} \delta_{xx'} \mathcal{J}_x^{ad}(l+l') , \\
[\mathcal{K}^{ab}(l), \mathcal{K}^{cd}(l')] & = \sqrt{G} \delta_{xx'} \sum_e \int_0^{l'} du \int_0^{l'-u} dv \{ \mathcal{J}_x^{eb}(l+l'-u-v) \Psi_x^{ad}(u) \Psi_x^{ce}(v) \\
& + \mathcal{J}_x^{ea*}(l+l'-u-v) \Psi_x^{de*}(u) \Psi_x^{cb}(v) \} \\
& - \sqrt{G} \delta_{xx'} \sum_e \int_0^l du \int_0^{l-u} dv \{ \mathcal{J}_x^{ed}(l+l'-u-v) \Psi_x^{cb}(u) \Psi_x^{ae}(v) \\
& + \mathcal{J}_x^{ec*}(l+l'-u-v) \Psi_x^{ad}(u) \Psi_x^{be*}(v) \} . \tag{13}
\end{aligned}$$

In the precise sense, these commutation relations are not an algebra because the open string creation operators $\Psi^{ab}(l)$ appear in the R.H.S. of (13). The algebraic structure (13) is understood as the consistency condition for the constraints on the equilibrium expectation value or “integrable condition” of the stochastic time evolution. The hermitian part of the current, $\mathcal{J}^{ab}(l) + \mathcal{J}^{*ba}(l)$, generates the S-D equation which implies the λ -independence of the equilibrium distribution. While the anti-hermitian part of the current, $\mathcal{J}^{ab}(l) - \mathcal{J}^{*ba}(l)$, satisfies $SU(r)$ current algebra. The similar algebraic structure is also found in Ref.[9] in a discretized form without continuum limit.

In conclusion, we have derived the continuum F-P hamiltonian which defines the non-critical orientable open-closed string field theory for $0 < c \leq 1$ by applying SQM to Kostov’s matrix-vector model. The origin of the algebraic structure (13) can be traced to the noise correlations which generate the time evolution in the Langevin equation. Namely, the noise correlations realize the deformation of open and closed strings which are equivalent to those generated by the three constraints, $\mathcal{L}(l)$, $\mathcal{J}^{ab}(l)$ and $\mathcal{K}^{ab}(l)$. We have shown that the structure to be universal representing the scale invariance in the non-critical orientable open-closed string theories. We hope that our approach is also usefull to the non-perturbative analysis of superstring theories such as type IIA, IIB and type I theories for which matrix models are proposed [14, 15] by the large N reduction of super Yang-Mills theory.

N. N. would like to thank I. Kostov for valuable comments. This work was partially supported by the Minisitry of Education, Science and Culture, Grant-in-Aid for Exploratory Research, 09874063, 1998.

References

- [1] E. Brézin and V. Kazakov, Phys. Lett. **B236**(1990) 144;
M. Douglas and S. Shenker, Nucl. Phys. **B335**(1990) 635;
D. Gross and A. Migdal, Phys. Rev. Lett. **64**(1990) 127; Nucl. Phys. **B340**(1990) 333.

- [2] N. Ishibashi and H. Kawai, Phys. Lett. **B314**(1993) 190;
Phys. Lett. **B322**(1994) 67.
- [3] A. Jevicki and J. Rodrigues, Nucl. Phys. **B421**(1994) 278.
- [4] M. Ikehara, N. Ishibashi, H. Kawai, T. Mogami, R. Nakayama and N. Sasakura,
Phys. Rev. **D50**(1994) 7467.
- [5] Y. Watabiki, Phys. Lett. **B346**(1995) 46;
Nucl. Phys. **B441**(1995) 119.
- [6] N. Nakazawa, Mod. Phys. Lett. **A10**(1995) 2175.
- [7] I. Kostov, Phys. Lett. **B344**(1995) 135;
Phys. Lett. **B349**(1995) 284.
- [8] T. Mogami, Phys. Lett. **B351**(1996) 439.
- [9] J. Avan and A. Jevicki, Nucl. Phys. **B469**(1996) 287.
- [10] N. Nakazawa and D. Ennyu, Phys. Lett. **B417**(1998)247.
- [11] I. Kostov, Nucl. Phys. **B376**(1992)539;
V. A. Kazakov and I. Kostov, Nucl. Phys. **B386**(1992)520.
- [12] V. A. Kazakov Phys. Lett. **B237**(1990) 212.
- [13] K. Ito, Proc. Imp. Acad. **20**(1944)519;
K. Ito and S. Watanabe, in Stochastic Differential Equations, ed. K. Ito (Wiley,
1978).
- [14] T. Banks, W. Fischler, S. Shenker and L. Susskind, Phys. Rev. **D55**(1997) 5112.
- [15] N. Ishibashi, H. Kawai, Y. Kitazawa and A. Tsuchiya, Nucl. Phys. **B498**(1997)469;
M. Fukuma, H. Kawai, Y. Kitazawa and A. Tsuchiya, hep-th/9705128.